

HSCRC Care Coordination:
The role of
Rehabilitation Services
February 12, 2015

Marlene Riley, MMS, OTR/L, CHT
Clinical Associate Professor
Towson University
College of Health Professions
Occupational Therapy Department

Key Ingredients & Program Elements of Successful Care Coordination Strategies

- Physical therapy, occupational therapy & speech-language pathology are integral services to address functional mobility, self-care, and cognitive function
- Rehabilitation Services are provided through full continuum of care: hospitalization , post-acute care (PAC) settings (SNF, In-pt rehab facility, LTC & HHA) & community-based
- Behavioral health occupational therapy across the continuum of inpatient, partial hospitalization, outpatient and home health
- Significant contribution to discharge plan and transitions between settings using most cost effective, evidence-based interventions

Barrier: Medicare B Therapy Cap

- Annual cap:

\$1,920 occupational therapy; up to \$3,700 with approved exception

\$1,920 physical therapy & SLP combined; up to \$3,700 with approved exception

- Includes: Observation days, SNF after first 21 days, LTC, out-patient therapy in hospital, free-standing out-patient or when non-homebound treated in home setting

- Maryland's rate-regulated environment impacts number of therapy visits for beneficiaries. Hospital out-pt may be most appropriate setting, but more costly than free-standing non rate-regulated setting. Variations in rate-regulated settings range from \$60 to \$225 for 30 minutes.

Additional Barriers

- Need for tool with standard patient assessment categories with distinct sensitivity for contributions of PT/ OT/ SLP for data collection across PAC settings
- Lack of data for costs associated with episodes of care across settings
- Lack of clear functional achievement goals at discharge with follow up by PAC services & impact to reduce unplanned readmissions
- Limited collaboration to connect individuals to community-based services after discharge from PAC settings

Workforce Requirements

- Multiple University Inter-professional Education Programs (PT, OT, SLP, PharmD, SW, RN)
- Multiple Campus-Community Partnerships
Ex. Gaitway 2 Better Balance with Baltimore Co Dept of Aging (BCDA)



Cockeysville Sr Center Fitness



State Senator Klausmeier, D-MD taking the Timed Up and Go test

Towson University Institute for Well-Being



HEARING AND BALANCE CENTER



[http://www.towson.edu/iwb/index.a
sp](http://www.towson.edu/iwb/index.asp)

Additional Reading

Andres, P.L., Haley, S.M., & Ni, P.S. (2003). Is patient-reported function reliable for monitoring post acute outcomes? *American Journal of Physical Medicine Rehabilitation*. 82(8), 614-621.

Bhattacharya, P., Khanal, D., Madhavan, R. & Chaturvedi, S. (2011). Why do ischemic stroke and transient ischemic attack patients get readmitted? *Journal of the Neurological Sciences*, 307, 50- 54.

Fisher, G. & Friesema. (2013). Implications of the affordable care act for occupational therapy practitioners providing services to medicare recipients. *The American Journal of Occupational Therapy*, 67 (5), 502 – 506.

Greysen, S.R., & Covinsky, K.E. (2014). Functional status—An important but overlooked variable in the readmissions equation. *Journal of Hospital Medicine*. 9 (5), 330 – 331.

Hoyer, E.H., Needham, D.M., Miller, J., Deutschendorf, A., Friedman, M., & Brotman, D.J. (2013). Functional status impairment is associated with unplanned readmissions. *Archives of Physical Medicine Rehabilitation*, 94 (10) , 1951-1958.

Jette, D. U., Stilphen, M., Ranganathan, V. K., Passek, S. D., Frost, F. S., & Jette, A. M. (2014). AM-PAC "6-Clicks" functional assessment scores predict acute care hospital discharge destination. *Physical Therapy* 94 (9), 1252 – 1261.

Krumholtz, H.M. (2013). Post-Hospital syndrome: An acquired, transient condition of generalized risk. *The New England Journal of Medicine*. 368 (2), 100 – 102.

Lord, R.K., Mayhew, C.R., Korupolu, R., Manthey, E.C., Friedman, M.A., Palmer, J.B., & Needham, D.M.(2013). ICU early physical rehabilitation programs: Financial modeling of cost savings. *Critical Care Medicine*. 41(3), 717-724.

Roberts, P.S. & Robinson, M.R. (2014). Occupational therapy's role in preventing acute hospital readmissions. *The American Journal of Occupational Therapy*, 68 (3), 1 -6.

Smith, B.A., Fields, C.J. & Fernandez, N. (2010). Physical therapists make accurate and appropriate discharge recommendations for patients who are acutely ill. *Physical Therapy* , 90 (5), 693 – 703.